

USGS Home

Contact USGS

Search USGS

Scientific Investigations Report 2006-5122

U.S. GEOLOGICAL SURVEY Scientific Investigations Report 2006-5122 DOE/ID-22198

A Conceptual Model of Ground-Water Flow in the Eastern Snake River Plain Aquifer at the Idaho National Laboratory and Vicinity with Implications for Contaminant Transport

Prepared in cooperation with the U.S. Department of Energy

By Daniel J. Ackerman, Gordon W. Rattray, Joseph P. Rousseau, Linda C. Davis, and Brennon R. Orr

Download the report (PDF, 6.56 MB)

Table of Contents

Conversion Factors and Datums

Abstract

Introduction

Conceptual Model of Ground-Water Flow

Implications for Contaminant Transport

Summary and Conclusions

Acknowledgments

References Cited

- •Appendix A. A Brief History of Wastewater-Disposal Practices at the Idaho National Laboratory
- •Appendix B. Levels of Information Needed for Constructing Conceptual Models at Different Scales

Figures

- Figure 1. Location of the Idaho National Laboratory, the model area, and selected facilities, wells, and streamflow-gaging stations.
- Figure 2. Distribution of selected waste contaminants in water from the eastern Snake River Plain aquifer, Idaho National Laboratory and vicinity, Idaho.
- Figure 3. Direction of regional ground-water flow, discharge areas, and irrigated acreage, eastern Snake River Plain, Idaho.
- Figure 4. Generalized geology, water-table altitude, and geologic groups and formations, and generalized stratigraphy of the eastern Snake River Plain, Idaho.
- Figure 5. Areas and grid sizes of ground-water models of the eastern Snake River Plain aquifer at the Idaho National Laboratory (INL) and vicinity, Idaho, relative to those of the Snake River Plain Regional Aquifer System Analysis (RASA) model.
- Figure 6. Composite stratigraphic units that form the eastern Snake River Plain aquifer, Idaho National Laboratory and vicinity, Idaho.
- Figure 7. Approximate locations of volcanic rift zones and the axial volcanic high,

Idaho National Laboratory and vicinity, Idaho.

- Figure 8. Locations of volcanic vents and vent corridors, Idaho National Laboratory and vicinity, Idaho.
- Figure 9. Distribution of hydrogeologic units present at the water table and water-table contours for 1980 in the model area, Idaho National Laboratory and vicinity, Idaho.
- Figure 10. Distribution of hydrogeologic units along the direction of regional ground-water flow in the model area, Idaho National Laboratory and vicinity, Idaho.
- Figure 11. Correlation among stratigraphic units of the eastern Snake River Plain, composite stratigraphic units beneath the Idaho National Laboratory, and hydrogeologic units of the conceptual model, Idaho National Laboratory and vicinity, Idaho.
- Figure 12. Water-table contours, 1995, and areas of abundant sediment within the model area, Idaho National Laboratory and vicinity, Idaho.
- Figure 13. Location of the model boundaries, Idaho National Laboratory and vicinity, Idaho.
- Figure 14. Depth to the base of the eastern Snake River Plain aquifer in the model area, Idaho National Laboratory and vicinity, Idaho.
- Figure 15. Contact between the intermediate-age rocks of hydrogeologic unit 3 and the older rocks underlying the base of the eastern Snake River Plain aquifer beneath the model area, Idaho National Laboratory and vicinity, Idaho.
- Figure 16. Regional Aquifer System Analysis model grid used to approximate the northeast, southeast-flowline, and southwest boundaries of the conceptual model, Idaho National Laboratory and vicinity, Idaho.
- Figure 17. Generalized ground-water budget components for the model area, 1980, Idaho National Laboratory and vicinity, Idaho.
- Figure 18. Mean annual streamflow at selected streamflow-gaging stations on the Big Lost River, Idaho, for 1965-2002.

- Figure 19. Distribution of lithium concentrations in water from wells in the eastern Snake River Plain aquifer, Idaho National Laboratory and vicinity, Idaho.
- Figure 20. Average linear ground-water velocities and flow directions calculated from the assumed first arrivals of chlorine-36 or the tritium/helium-3- and chlorofluorocarbon-model ages of water from selected wells, Idaho National Laboratory and vicinity, Idaho.
- Figure 21. Temperature of ground water in samples from wells, Idaho National Laboratory and vicinity, Idaho.
- Figure 22. Generalized water-level changes at the Idaho National Laboratory and vicinity, Idaho, for July 1981–July 1985, April–August 1988 to April–August 1991, March–May 1991 to March–May 1995, and March–May 1995 to March–May 1998.
- Figure 23. Water levels in wells Arbor Test, USGS 9, USGS 12, and USGS 25 for 1948–2003, Idaho National Laboratory, Idaho.
- Figure 24. Distribution of hydrogeologic units and the direction of ground-water flow for the model area, Idaho National Laboratory and vicinity, Idaho.

Tables

- Table 1. Summary of surface and subsurface wastewater disposal at selected facilities, Idaho National Laboratory, Idaho.
- Table 2. Ranges of hydraulic conductivities and porosities for hydrogeologic units and other rocks of hydrogeologic importance, Idaho National Laboratory and vicinity, Idaho.
- Table 3. Summary statistics of basalt flow thickness within individual hydrogeologic units, Idaho National Laboratory and vicinity, Idaho.
- Table 4. Summary of conceptual model boundary characteristics and estimated 1980 inflows and outflows, Idaho National Laboratory and vicinity, Idaho.
- Table 5. Summary of streamflow at two streamflow-gaging stations on the Big Lost River, Idaho.
- Table 6. Ground-water budget for the conceptual model area for 1980 and

derived from the Regional Aquifer System Analysis (RASA) model of the Snake River Plain aquifer, Idaho.

This report is available online in Portable Document Format (PDF). If you do not have the **Adobe Acrobat PDF Reader**, it is available for free download from Adobe Systems Incorporated.

Download the report (PDF, 6.56 MB)

Document Accessibility: Adobe Systems Incorporated has information about PDFs and the visually impaired. This information provides tools to help make PDF files accessible. These tools convert Adobe PDF documents into HTML or ASCII text, which then can be read by a number of common screen-reading programs that synthesize text as audible speech. In addition, an accessible version of Acrobat Reader 7.0 for Windows (English only), which contains support for screen readers, is available. These tools and the accessible reader may be obtained free from Adobe at Adobe Access.

Send questions or comments about this report to the author, D.J. Ackerman, (208) 526-2062.

For more information about USGS activities in Idaho, visit the USGS Idaho Water Science Center home page.

AccessibilityFOIAPrivacyPolicies and Notices

U.S. Department of the Interior | U.S. Geological Survey Persistent URL: http://pubs.water.usgs.gov/sir20065122

Page Contact Information: Publications Team

Page Last Modified: Thursday, 24-Aug-2006 14:44:27 EDT

